

Thimerosal Content in Some US Licensed Vaccines
updated October 2004

Vaccine	Brand Name	Manufacturer	Thimerosal Concentration(1)	Mercury ug/0.5 ml	
Anthrax	Biothrax	BioPort Corp	0	0	
	Tripedia	Aventis Pasteur	*	*	
DTaP	Infanrix	GlaxoSmithKline	0	0	
	Daptacel	Aventis Pasteur	0	0	
DTaP - Hep B - IPV	Pediarix	GlaxoSmithKline	*	*	
DTwP	All Products		.01%	25	
DT	Diphtheria & Tetanus Toxoids Adsorbed USP	multi-dose	Aventis Pasteur	.01%	25
		single dose		*	*
Td	Tetanus and Diphtheria Toxoids Adsorbed	Aventis Pasteur	.01%	25	
Tetanus Toxoid	Tetanus Toxoid Adsorbed USP		.01%	25	
	Tetanus Toxoid Adsorbed Adult Use	Aventis Pasteur	.01%	25	
	Booster		.01%	25	
Hib	ActHIB /OmniHib	Aventis Pasteur	0	0	
	TriHIBit	Aventis Pasteur	*	*	
	HibTITER	Wyeth-Ayerst	0	0	
Hib/HepB	PedvaxHIB liquid (2)	Merck	0	0	
	Comvax (3)	Merck	0	0	
Hepatitis A	Havrix	GlaxoSmithKline	0	0	
	Vaqta adult/pediatric	Merck	0	0	
Hepatitis B	Engerix-B preservative free	GlaxoSmithKline	*	*	
	Recombivax HB preservative free	Merck	0	0	
Hep A-B	Twinrix	GlaxoSmithKline	*	*	
			*	*	
Influenza 2004 /5 Formula	Fluzone	prefilled syringe (0.25 mL)			
		prefilled syringe (0.5mL)	Aventis Pasteur	.01%	25
		multi-dose (5 mL)		.01%	25
	Fluvirin preservative free	Evans	*	*	
IPV	Fluvirin		.01%	25	
	FluMist	MedImmune	0	0	
Meningococcal	MENOMUNE-A/C/Y/W-135	multi-dose	Aventis Pasteur	.01%	25
		single dose		*	*
MMR	M-M-R II	Merck	0	0	
Pneumococcal	Prennar	Wyeth-Ayerst	0	0	
	Pneumovax 23	Merck	0	0	
Rabies	RabAvert	Chiron	0	0	
	IMOVAX	Aventis Pasteur	0	0	
Typhoid Fever	Typhim VI	Aventis Pasteur	0	0	
	Vivotif	Berna Biotch	0	0	
Varicella	Varivax	Merck	0	0	
Yellow Fever	YF-VAX	Aventis Pasteur	0	0	

1. A concentration of 1:10,000 is equivalent to a 0.01% concentration. Thimerosal is approximately 50% Hg by weight. A 1:10,000 concentration contains 25 micrograms of Hg per 0.5 mL.

2. A previously marketed lyophilized preparation contained 0.005% thimerosal.

3. COMVAX is not approved for use under 6 weeks of age because of decreased response to the Hib component.

* This product should be considered equivalent to thimerosal-free products. This vaccine may contain trace amounts (<0.3 µg) of mercury left after post-production thimerosal removal; these amounts have no biological effect. JAMA 1999;282(18) and JAMA 2000;283(16).

Institute for Vaccine Safety



Johns Hopkins Bloomberg
School of Public Health

www.vaccinesafety.edu